Surname

Centre Number

Other Names

wjec

### GCE AS/A LEVEL

2410U10-1

### CHEMISTRY – AS unit 1 The Language of Chemistry, Structure of Matter and Simple Reactions

TUESDAY, 22 MAY 2018 - MORNING

1 hour 30 minutes

	For Exa	e only	
	Question	Maximum Mark	Mark Awarded
Section A	1. to 6.	10	
Section B	7.	12	
	8.	15	
	9.	15	
ed a:	10.	13	
	11.	15	
	Total	80	

#### ADDITIONAL MATERIALS

In addition to this examination paper, you will need a:

- · calculator;
- Data Booklet supplied by WJEC.

#### INSTRUCTIONS TO CANDIDATES

Use black ink or black ball-point pen. Do not use gel pen or correction fluid.

Write your name, centre number and candidate number in the spaces at the top of this page.

**Section A** Answer **all** questions in the spaces provided.

Section B Answer all questions in the spaces provided.

Candidates are advised to allocate their time appropriately between **Section A (10 marks)** and **Section B (70 marks)**.

#### **INFORMATION FOR CANDIDATES**

The number of marks is given in brackets at the end of each question or part-question.

The maximum mark for this paper is 80.

Your answers must be relevant and must make full use of the information given to be awarded full marks for a question.

The assessment of the quality of extended response (QER) will take place in Q.11(a)(i).

If you run out of space, use the additional page(s) at the back of the booklet, taking care to number the question(s) correctly.



		SECTI	ON A		
	Answ	ver <b>all</b> questions in	the spaces pro	ovided.	
For the ionic compound caesium chloride, state the coordination number of the chloride ion. [1]					
The first fou	ur ionisation ener	rgies for an eleme	nt are shown in	the table below.	
		Ionisation ener	rgy / kJ mol <sup>-1</sup>		
	1st	2nd	3rd	4th	
	548	1060	4120	5440	
	g allowe to ropi				re of a chromium
atom.	3s	Зр	30		[1] 4s
neon core	3s				
neon	3s				
neon	3s				
neon	3s				
neon	3s				
neon	3s				
neon	3s				
neon	3s				




4.		sphorus(V) chloride reacts with water to form phosphoric acid and hydrogen chloride reactants and products are shown in the equation below but the equation is <b>not</b> balanced.	
		PCI <sub>5</sub> + H <sub>2</sub> O → H <sub>3</sub> PO <sub>4</sub> + HCI	
	(a)	Balance the equation. [1]	
	(b)	Calculate the atom economy for the formation of phosphoric acid in this reaction. [2]	
		Atom economy =	
5.	(a)	Give the meaning of the term <i>electronegativity</i> . [1]	2410U101 03
	(b)	Explain why electronegativity increases across a period in the Periodic Table. [1]	
6.	Brom (a)	nine is produced commercially from the bromide ions in sea water by reaction with chlorine Give the <b>ionic</b> equation for this reaction.	
	(b)	Describe a test, apart from the use of chlorine, to show that a solution contains bromide	
		ions. Give the reagent(s) and observation(s). [1] Reagent(s)	
		Observation(s)	
			10



	SECTION B
	Answer all questions in the spaces provided.
(a)	Melting temperatures vary down groups and across periods.
	<ul> <li>(i) Explain why chlorine is a gas but iodine is a solid at room temperature and pressure.</li> <li>[3]</li> </ul>
	(ii) Explain why sodium has a lower melting temperature than aluminium. [1
	(iii) Explain why silicon has a higher melting temperature than phosphorus. [1
(b)	State and explain how you would expect the first ionisation energy of nitrogen to compare with the first ionisation energy of oxygen.
······	
•••••	



© WJEC CBAC Ltd.



© WJEC CBAC Ltd.

(2410U10-1)

Turn over.

	gives soi	me information about four r	auloactive isotopes.	
Isotope	е	Radiation emitted	Half-life	
<sup>90</sup> Sr		β	28 years	
<sup>99</sup> Tc		γ	6 hours	
<sup>210</sup> At		α	8.1 hours	
<sup>228</sup> Th		α	1.9 years	



# **BLANK PAGE**

7

## **PLEASE DO NOT WRITE ON THIS PAGE**





(a)	The diagram below shows part of the atomic emission spectrum of hydrogen.					
		increasing frequency —				
	(i)	Use the letter <b>A</b> to label the line of longest wavelength on the diagram. [1				
	(ii)	Explain why hydrogen atoms emit only certain definite frequencies of visible light. [2				
	······					
(b)	The (i)	ionisation energy of a hydrogen atom is $2.18 \times 10^{-21}$ kJ. Explain what this statement means. [2				
	(ii)	Calculate the minimum frequency of radiation required to ionise a hydrogen atom i its ground state.				
		Frequency = s <sup>-</sup>				



		Examiner
(C)	Hydrazine is a compound of hydrogen and nitrogen only. It is a colourless, flammable liquid which was used in various rocket fuels.	only
	0.160 g of hydrazine on vaporisation at 398 K and 1 atm pressure has a volume of $163  \text{cm}^3$ .	
	Calculate its volume at 273 K and 1 atm pressure and hence show that its molecular formula is $N_2H_4$ . [3]	
••••••		
(d)	(i) Draw a dot and cross diagram to show the electron arrangement in hydrazine, $N_2H_4$ . Show outer electrons only. [2]	101
		2410U101
	(ii) Hydrazine contains polar covalent bonds between nitrogen and hydrogen atoms.	
	State what is meant by a <i>polar</i> covalent bond. [1]	
(e)	Hydrazine acts as a base in a similar way to ammonia.	
	Suggest an equation for the equilibrium formed when hydrazine dissolves in water. [1]	
		15

(2410U10-1)

© WJEC CBAC Ltd.



Turn over.

2410U101 09

Ascorbic acid, C <sub>6</sub> H <sub>8</sub> O <sub>6</sub> , is the main component of vitamin C tablets. Its name is derived from a (meaning "no") and <i>scorbutus</i> (scurvy), the disease caused by a deficiency of vitamin C. A student was asked to find the percentage of ascorbic acid in identical vitamin C tablets. She was told to use the following method.  Fill a burette with 0.100 mol dm <sup>-3</sup> sodium hydroxide solution. Weigh a conical flask and record its mass. Add a vitamin C tablet to the flask, reweigh it and record its mass. Add a bout 50 cm <sup>3</sup> of deionised water to the flask and swirl to break up the tablet. Heat the flask gently for 5 to 10 minutes. After the solution has cooled add a few drops of a suitable indicator. Accurately repeat the procedure several times and calculate a mean titre. (a) A three decimal place balance was used. The mass of each vitamin C tablet was 500 mg. Calculate the maximum percentage error in the weighing of the tablet. You must show your working. (b) (i) Suggest why she did not need to measure the volume of water accurately. (1] (ii) Suggest why she heated the flask for 5 to 10 minutes. [1]	Asco		
<ul> <li>Fill a burette with 0.100 dm<sup>-3</sup> sodium hydroxide solution.</li> <li>Weigh a conical flask and record its mass.</li> <li>Add a vitamin C tablet to the flask, reweigh it and record its mass.</li> <li>Add about 50 cm<sup>3</sup> of deionised water to the flask and swirl to break up the tablet.</li> <li>Heat the flask gently for 5 to 10 minutes.</li> <li>After the solution has cooled add a few drops of a suitable indicator.</li> <li>Carry out a rough titration of this solution with the sodium hydroxide solution.</li> <li>Accurately repeat the procedure several times and calculate a mean titre.</li> </ul> (a) A three decimal place balance was used. The mass of each vitamin C tablet was 500 mg. Calculate the maximum percentage error in the weighing of the tablet. You must show your working. [2] Maximum percentage error =% (b) (i) Suggest why she did not need to measure the volume of water accurately. [1]	<i>a</i> - (n	eaning "no") and scorbutus (scurvy), the disease caused by a deficiency of vitami	
<ul> <li>Weigh a conical flask and record its mass.</li> <li>Add a vitamin C tablet to the flask, reweigh it and record its mass.</li> <li>Add about 50 cm<sup>3</sup> of deionised water to the flask and swirl to break up the tablet.</li> <li>Heat the flask gently for 5 to 10 minutes.</li> <li>After the solution has cooled add a few drops of a suitable indicator.</li> <li>Carry out a rough titration of this solution with the sodium hydroxide solution.</li> <li>Accurately repeat the procedure several times and calculate a mean titre.</li> <li>(a) A three decimal place balance was used. The mass of each vitamin C tablet was 500 mg. Calculate the maximum percentage error in the weighing of the tablet. You must show your working.</li> <li>(a) Maximum percentage error =%</li> <li>(b) (i) Suggest why she did not need to measure the volume of water accurately. [1]</li> </ul>	She	was told to use the following method.	
Calculate the maximum percentage error in the weighing of the tablet.       [2]         You must show your working.       [2]         Maximum percentage error =	• • • •	<ul> <li>Weigh a conical flask and record its mass.</li> <li>Add a vitamin C tablet to the flask, reweigh it and record its mass.</li> <li>Add about 50 cm<sup>3</sup> of deionised water to the flask and swirl to break up the tablet.</li> <li>Heat the flask gently for 5 to 10 minutes.</li> <li>After the solution has cooled add a few drops of a suitable indicator.</li> <li>Carry out a rough titration of this solution with the sodium hydroxide solution.</li> </ul>	
You <b>must</b> show your working. [2] Maximum percentage error =	(a)	A three decimal place balance was used. The mass of each vitamin C tablet was 500	) mg.
(b) (i) Suggest why she did not need to measure the volume of water accurately. [1]			[2]
(ii) Suggest why she heated the flask for 5 to 10 minutes. [1]	(b)		
		(ii) Suggest why she heated the flask for 5 to 10 minutes.	[1]



Examiner only The student used the results from three titrations to calculate a mean titre. Some of her (C) results are shown below. Titration 1 2 3 Final reading / cm<sup>3</sup> 26.90 26.90 Initial reading / cm<sup>3</sup> 0.25 0.20 0.15 Titre / cm<sup>3</sup> 26.65 26.75 Mean titre =  $26.73 \text{ cm}^3$ Determine the **final reading** for the third titration. [2] Final reading = ..... cm<sup>3</sup> Ascorbic acid can decompose upon exposure to air. If this reaction occurred before (d) the titration was completed, state how it might affect the titration results. Explain your answer. [2]



Examiner only The equation for the reaction between ascorbic acid and sodium hydroxide is given below. (e)  $C_6H_8O_6$  + NaOH  $\longrightarrow$   $C_6H_7O_6Na$  +  $H_2O$ *M*<sub>r</sub> 176 The percentage of ascorbic acid is identical in each 500 mg tablet. Calculate the percentage of ascorbic acid in each vitamin C tablet. [3] Percentage ascorbic acid = ......% Sulfuric acid and hydrochloric acid are strong acids. (**f**) (i) Calculate the pH of a solution of 0.010 mol dm<sup>-3</sup> sulfuric acid, H<sub>2</sub>SO<sub>4</sub>. [2] pH = ..... When hydrochloric acid is heated with  ${\rm MnO}_2$  it reacts according to the following (ii) equation.  $MnO_2(s) + 4HCl(aq) \longrightarrow MnCl_2(aq) + Cl_2(g) + 2H_2O(l)$ Explain why this can be classified as a redox reaction. [2] 15



# **BLANK PAGE**

13

## PLEASE DO NOT WRITE ON THIS PAGE



(a)	Nitro	ogen can react with hydrogen to form ammonia.	E>
		$N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$	
	A mi The	xture of $N_2$ and $H_2$ is left to react at a certain temperature, until it reaches equilibrium equilibrium mixture has the following composition.	ı.
		$\begin{array}{ll} N_2 & 1.16  \text{mol}  \text{dm}^{-3} \\ H_2 & 1.60  \text{mol}  \text{dm}^{-3} \\ NH_3 & 0.752  \text{mol}  \text{dm}^{-3} \end{array}$	
	(i)	A student said that the equilibrium must lie to the left because the concentrations on nitrogen and hydrogen are greater than that of ammonia. Is he correct?	of
		Justify your answer by calculating a value for $K_c$ for this equilibrium. Give the unit for $K_c$ . [4	.]
		K <sub>c</sub> =	
		Unit	
	·····		
	(ii)	When the temperature is increased the equilibrium yield of $NH_3$ decreases. The student said that the reaction is endothermic. Is he correct?	e
		Justify your answer by using Le Chatelier's principle. [2]	2]



Examiner only Ammonia can be used as part of the nitrophosphate process to produce the fertiliser diammonium hydrogenphosphate (DAP) which has the formula  $(NH_4)_2HPO_4$ . (b)  $Ca(NO_3)_2 + 4H_3PO_4 + 8NH_3 \longrightarrow CaHPO_4 + 2NH_4NO_3 + 3(NH_4)_2HPO_4$ *M*<sub>r</sub> 132 Calculate the maximum mass of DAP, in kg, that could be made from 1.00 tonne of ammonia. [3] Maximum mass = ...... kg Calculate the volume, in cm<sup>3</sup>, that  $2.54 \times 10^{-3}$  mol of nitrogen occupies at a temperature (C) of 120 °C and a pressure of 101 kPa. [4] Volume = ..... cm<sup>3</sup> 13



			٦Ex
. <i>(a)</i>	A so prese	plution contains one cation and up to three different anions. The anions possibly ent are carbonate, chloride and sulfate.	
	(i)	Devise a plan that unambiguously proves which anions are present in the mixture.	
		You should also give any observations and conclusions that enable you to identify the anions. [6 QER]	
	·····		
	•••••		
	•••••		
	•••••		
	•••••		
	•••••		
	•••••		



(ii) A student said that the cation in the mixture can only be sodium since all the possible anions form a soluble salt with sodium. Is he correct?

Justify your answer and state how you could prove if the statement were true. [2]

(b) A student is given four materials and asked to identify the **type of structure** present in each one by carrying out a series of tests.

She is told that the maximum temperature of a Bunsen burner flame is about 800 °C. She is also told that in at least one case, it will **not** be possible to come to a definite conclusion.

Her results are shown in the table below.

	Α	В	С	D
Melting temperature / °C	100	>800	>800	>800
Solubility in water	soluble	insoluble	insoluble	soluble
Conductivity of solid	none	none	good	none
Conductivity of solution	none			good

(i) Use the information in the table to identify each type of structure. Where a definite conclusion cannot be reached explain your reasoning.

[4]

Examiner only

**QUESTION CONTINUES ON PAGE 18** 



Examiner only For **one** of the materials where the type of structure could not be identified, suggest what further test(s) are needed to identify the type of structure. [2] (ii) Suggest why it is difficult to identify a material as a metal when it is in powdered (iii) form. [1] 15 **END OF PAPER** 

© WJEC CBAC Ltd.

(2410U10-1)

# **BLANK PAGE**

19

## PLEASE DO NOT WRITE ON THIS PAGE



Question number	Additional page, if required. Write the question number(s) in the left-hand margin.	Examine only

